

Redox-Active Ligands

Redox-Active Ligand-Induced Homolytic Bond Activation**

Daniël L. J. Broere, Lotte L. Metz, Bas de Bruin, Joost N. H. Reek, Maxime A. Siegler, and Jarl Ivar van der Vlugt*

Dedicated to Professor Thomas B. Rauchfuss on the occasion of his 65th birthday


Abstract: Coordination of the novel redox-active phosphine-appended aminophenol pincer ligand (PNO^{H2}) to Pd^{II} generates a paramagnetic complex with a persistent ligand-centered radical. The complex undergoes fully reversible single-electron oxidation and reduction. Homolytic bond activation of diphenyldisulfide by the single-electron reduced species leads to a ligand-based mixed-valent dinuclear palladium complex with a single bridging thiolate ligand. Mechanistic investigations support an unprecedented intramolecular ligand-to-disulfide single-electron transfer process to induce homolytic S–S cleavage, thereby releasing a thiyl (sulfanyl) radical. This could be a new strategy for small-molecule bond activation.

Redox-active ligands are frequently encountered in important natural processes mediated by metalloenzymes.^[1] In inorganic chemistry, these systems have long been considered to be primarily a spectroscopic curiosity, with major focus on understanding the electronic structure and bonding within homoleptic systems.^[2] Recently, heteroleptic complexes have been shown to offer unique reactivity in stoichiometric activation reactions and in catalysis, since the redox-active nature of these ligands allows their use as an electron reservoir during (catalytic) turnover.^[3] The majority of redox-active frameworks are based on nitrogen or oxygen donors,^[4] with aminophenol-based *N,O* ligands as archetypical redox-active systems that can span three oxidation states.^[5] In contrast and at odds with the relevance of phosphorus ligands in homogeneous catalysis, few phosphine-containing redox-active ligands exist.^[6,7] Thomas and co-workers recently described an (*o*-anilino)phenylphosphine ligand that is susceptible to oxidation in the coordination sphere of Cu^I, but radical P–P coupling precluded the use of

this scaffold as a reversible redox-active ligand.^[8] Installment as a redox-innocent entity adjacent to a redox-active framework is less likely to affect the coordinative properties at phosphorus, but relatively few of these ligands have been developed.^[9,10]

Bond homolysis is a very useful reaction to probe for accessible ligand-based reactivity. Established ligand-mediated bond activation (and formation) reactions classify overall as two-electron processes.^[11–14] Metal-mediated one-electron homolysis is much rarer,^[15] while reductive homolytic bond fission originating from ligand-based overall single-electron transfer is, to the best of our knowledge, unknown (Figure 1). Methodologies that facilitate odd-electron transfer processes will allow the controlled generation of reactive substrate radicals for synthetic chemistry.^[16]

Common: two-electron homolysis of e.g. disulfides



M = noble metal

Unknown: ligand-induced reductive one-electron bond homolysis




Figure 1. a) Typical reactivity concerning bond homolysis by noble metal complexes. b) Unprecedented reductive single-electron transfer from a redox-active ligand to a disulfide substrate, generating a thiolate and a thiyl radical.

Recently, we reported a tridentate redox-active NNO ligand that accommodates radical-type C–H amination reactivity on a Pd^{II} platform.^[17] In order to arrive at a redox-active phosphine ligand, we sought to merge the redox-active aminophenol framework with a flanking diphenylphosphine group. Addition of this (sterically encumbered) donor should impact the redox properties of the *N,O* moiety upon coordination to a transition metal, relative to the previous NNO scaffold. We herein describe the facile synthesis and electronic structure of a phosphorus ligand that is ‘redox active’ when coordinated to Pd^{II}. This system, which displays a markedly lower reduction potential than the Pd complex with our previously reported NNO system,^[17] is able to facilitate radical-type homolytic bond activation of disulfides, with formation of a well-defined ligand-based mixed-valent dinuclear complex.


The novel aminophenolphosphine ligand PNO^{H2} (³¹P NMR $\delta = -20.25$) was prepared as an air-sensitive

[*] D. L. J. Broere, L. L. Metz, Prof. Dr. B. de Bruin, Prof. Dr. J. N. H. Reek, Dr. J. I. van der Vlugt
Homogeneous, Bioinspired & Supramolecular Catalysis
van't Hoff Institute for Molecular Sciences
University of Amsterdam
Science Park 904, 1098 XH Amsterdam (The Netherlands)
E-mail: j.i.vandervlugt@uva.nl

Dr. M. A. Siegler
Department of Chemistry, John Hopkins University
3400 N Charles Street, Baltimore, MD 21218 (USA)

[**] Research funded by the European Research Council (ERC Starting Grant 279097 to J.I.v.d.V.). We thank Ed Zuidinga for assistance with ESI-MS measurements.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201410048>.



Scheme 1. Synthetic route to complexes **1** and **2**.

white solid in 58% overall yield through a two-step procedure from commercially available *o*-iodoaniline and 3,5-di-*tert*-butylcatechol.^[18] Reaction with $[\text{PdCl}_2(\text{MeCN})_2]$ gave complex **1** as an orange solid in 67% yield (^{31}P NMR $\delta = 43.98$). Addition of triethylamine resulted in a rapid color change to green, and subsequent exposure to air afforded the dark-red paramagnetic species **2** in 78% yield (Scheme 1). Magnetic susceptibility measurements (Evans' method) showed an effective magnetic moment (μ_{eff}) of $1.81 \mu_{\text{B}}$, thus indicating an $S = 1/2$ ground state. Hence, this species is best formulated as $[\text{PdCl}(\text{PNO}^{\text{ISQ}})]$, and this assignment was confirmed by single-crystal X-ray diffraction (Figure 2; ISQ = iminosemiquinonato).




Figure 2. a) Displacement ellipsoid plot (50% probability level) of complex **2** at 110(2) K. b) Relevant experimental (XRD) and computed (DFT) metric parameters support the PNO^{ISQ} state in **2**. Selected bond angles [°]: $\text{N}_1\text{-Pd}_1\text{-Cl}_1$ 174.90(5); $\text{P}_1\text{-Pd}_1\text{-O}_1$ 167.43(4); $\text{N}_1\text{-Pd}_1\text{-Cl}_1$ 94.038(17); $\text{N}_1\text{-Pd}_1\text{-O}_1$ 81.16(6).

Complex **2** shows a slightly distorted square-planar geometry with an acute $\angle \text{N}_1\text{-Pd}_1\text{-O}_1$ angle of $81.16(6)^\circ$. Palladium–ligand bond lengths and angles in **2** compare well with PdCl complexes bearing redox-innocent monoanionic PNO pinners.^[19] The metric parameters found for the amidophenolate fragment support the ISQ oxidation state of the ligand^[20] and these data are reproduced by DFT (b3-lyp/def2-TZVP) optimized geometric parameters for the doublet PNO^{ISQ} ground state. X-band EPR spectroscopy in toluene at 298 K revealed hyperfine couplings with ^{105}Pd , ^{31}P , ^{14}N , and three ^1H nuclei (see Table S1 in the Supporting Information). The g_{iso} value of 2.0052 suggests coordination of a PNO^{ISQ} ligand radical to Pd^{II} .^[21] The calculated spin density for **2** predominantly resides on the ligand (97% total spin density, 21% on the iminosemiquinonato nitrogen), in agreement with EPR observations (Figure 3 and the Supporting Information).

Cyclic voltammetry of **2** in CH_2Cl_2 solution shows fully reversible one-electron oxidation and reduction events at





Figure 3. a) Experimental and simulated EPR spectrum of **2** (toluene, RT) Freq = 9.366829 GHz, $T = 298$ K, Mod Ampl. = 1 Gauss, power = 20 mW. Simulated (DFT) g value and hyperfine couplings A (MHz): g_{iso} 2.0052 (2.0062); A^{Pd} +10.70 (+8.61); A^{P} -13.55 (-16.04); A^{N} +16.45 (+11.0); A^{H^1} -4.90 (-5.06); A^{H^2} (NR) (-0.78); A^{H^3} -4.95 (-5.24); A^{H^4} (NR) (2.60); A^{H^5} -7.18 (-5.66); A^{H^6} (NR) (2.15) NR = not resolved; DFT parameters: ORCA (b3-lyp, def2-TZVP). b) DFT (b3-lyp/def2-TZVP) calculated spin-density plot for **2**.

+0.07 V and -0.75 V vs. Fc/Fc^+ , respectively. Reversible ligand-based redox chemistry was also observed by UV/Vis spectroelectrochemistry in an optically transparent thin-layer electrolysis (OTTLE) cell (see the Supporting Information for details of the redox chemistry). Chemical reduction of **2** with $[\text{CoCp}_2]$ in CH_2Cl_2 furnished air-sensitive diamagnetic complex **3**, formulated as $[\text{CoCp}_2][\text{PdCl}(\text{PNO}^{\text{AP}})]$ (^{31}P NMR $\delta = 36.56$; AP = amidophenolato). To assess the steric constraint imposed on the Pd center by the flanking phosphine donor, complex **3** was exposed to exogenous phosphines. No reaction was observed with PPh_3 but coordination of PMe_3 rapidly produced complex **4** (^{31}P NMR $\delta = 41.52$ (d) and -10.61 (d); $J_{\text{P-P}}$ 40.2 Hz), formulated as $[\text{Pd}(\text{PMe}_3)(\text{PNO}^{\text{AP}})]$ (Scheme 2).



Scheme 2. Synthetic route to complexes **3** and **4**.

Oxidative addition of a disulfide to low-valent Pd is usually a two-electron process.^[22] Given the demonstrated reversible one-electron chemistry of species **3** at a mild potential, we sought to investigate its reactivity toward disulfides. Addition of TIPF_6 to a suspension of **3** in benzene in the presence of an equimolar amount of diphenyldisulfide produced the soluble paramagnetic species **5**. Magnetic susceptibility measurements of **5** at 298 K using Evans' method gave an effective magnetic moment (μ_{eff}) of $1.90 \mu_{\text{B}}$, thus indicating an $S = 1/2$ ground state. This observation implies one-electron oxidation of PNO^{AP} to PNO^{ISQ} . CSI-MS studies in benzene indicate the presence of a dinuclear species in solution at m/z 1279.28 $[M]^+$, formulated as $[\text{Pd}_2(\mu$

SPh)(PNO)₂]. UV/Vis spectroscopy shows characteristic absorption bands for both PNO^{AP} and PNO^{ISO} ligand fragments. X-band EPR spectroscopy of compound **5** in toluene at 298 K showed an isotropic signal with no resolved hyperfine couplings. The *g*_{iso} value of 2.0041 supports the presence of a PNO^{ISO} ligand radical. This assignment was corroborated by X-ray diffraction (Figure 4).





Figure 4. a) Displacement ellipsoid plot (50% probability level) of complex **5** at 110(2) K. Disorder in the 3,5-*t*Bu₂Ph ring at Pd₂ and disordered solvent omitted for clarity. b) metric parameters for the PNO^{ISO} and PNO^{AP} fragments in **5**. Selected additional bond lengths [Å] and angles [°]: Pd1–N11 1.990(3); Pd2–N12 2.011(3); Pd1–O11 2.041(3); Pd2–O12 2.063(3); Pd1–S1 2.3203(9); Pd2–S1 2.3012(10); Pd1–S1–Pd2 104.58(4); N11–Pd1–S1 178.45(10); N12–Pd2–S1 175.5(4); N11–Pd1–O11 82.63(12); N12–Pd2–O12 81.5(3). Atom numbering: first digit = atom number, second digit = ligand number.

The molecular structure contains one thiophenolate unit bridging two Pd^{II}(PNO) centers. This bridging monothiolate motif, although not unique, is rather uncommon, particularly with Pd.^[23] Strikingly, the observed metric parameters indicate different oxidation states for the two PNO ligands present, that is, the amidophenolato (N11–O11) and the iminosemiquinonato (N12–O12) forms. VT-EPR spectroscopic data indicate facile electron exchange between the PNO^{AP} and the PNO^{ISO} moieties of **5** in solution (see the Supporting Information). We are not aware of similar examples of monobridged dinuclear complexes that show ligand-based mixed valency.^[24] For homodinuclear reaction centers, the mixed valency is typically metal-centered or shared between the metal and (bridging) ligand.^[25] Systems with separated mixed-valent ligand-based redox centers could be of interest for studying intramolecular electron-transfer processes and potentially also for ligand-

assisted redox catalysis.^[26] Selective formation of such species through controllable synthetic procedures from stable monomeric precursors might allow the study of these electronic configurations.

The occurrence of outer-sphere electron transfer from **2** to PhSSPh is excluded on the basis of their relative redox potentials.^[26] The formation of species **5** (Scheme 3) is proposed to involve initial chloride dissociation and disulfide coordination. Dialkyl disulfides have a higher S–S bond dissociation energy than diaryl disulfides and are thus less prone to undergoing bond homolysis.^[27] Using di(*tert*-butyl)disulfide instead of PhSSPh allowed observation of the corresponding Pd–disulfide adduct by NMR spectroscopy. The ³¹P NMR chemical shift of $\delta = 39.82$ is similar to that of neutral **4**. The non-equivalent *tert*-butyl groups of the substrate are shifted upfield in the ¹H NMR spectrum, which otherwise resembles that of **4** (see the Supporting Information). Subsequent intramolecular ligand-to-substrate single-electron transfer results in homolytic S–S bond cleavage with formation of [PNO^{ISO}PdSPh] and release of a PhS[•] radical. This thiyl radical can either undergo self-recombination or react with a ‘vacant’ [PNO^{AP}Pd] complex, thus forming PhSSPh or a second equivalent of [PNO^{ISO}PdSPh], respectively. The final step is the formation of the mixed-valent [(PNO^{ISO})Pd(μ-SPh)Pd(PNO^{AP})] (**5**), through coordination of a sulfur lone pair in [PNO^{ISO}PdSPh] to free [PNO^{AP}Pd]. Starting with a 4:1 ratio of **3**/PhSSPh also leads to the clean production of complex **5**, thus supporting this pathway. Detection of thiyl radicals by EPR spectroscopy with DMPO (DMPO = 5,5-dimethyl-1-pyrroline *N*-oxide) as a spin-trapping agent was unsuccessful, probably owing to a high recombination rate relative to the generation rate of these thiyl radicals, the short lifetime of DMPO·(SPh) adducts,^[28] and the competitive reaction of complex **3** with DMPO. However, GC–MS analysis of the reaction mixture confirmed the presence of diphenylsulfide, which is generated from the reaction of PhS[•] with the solvent benzene. Using a mixture of PhSSPh and di(*p*-tolyl)disulfide



Scheme 3. Proposed mechanism for the formation of dinuclear [(PNO^{ISO})Pd(μ-SPh)Pd(PNO^{AP})] (**5**) with mixed valency in the two PNO scaffolds.

led to co-formation of phenyl(*p*-tolyl)disulfide, as detected by GC-MS, thus supporting the intermediacy of thiyl radicals created by this ligand-to-substrate electron transfer process.

In conclusion, the first example of a phosphine ligand appended to a redox-active aminophenol framework is reported. This PNO^{H2} pincer ligand can coordinate to Pd^{II} as a neutral (**1**), radical monoanionic (**2**), or dianionic scaffold (**3,4**), as supported by spectroscopic, X-ray crystallography, and computational data. Cyclic voltammetry and spectroelectrochemistry demonstrate reversible single-electron redox events for complex **2**. The bulky phosphine arm and rigid backbone enforce considerable steric crowding around the Pd center. One-electron reduction generates complex **3**, which is a competent reagent for homolytic bond activation of disulfides through ligand-to-substrate single-electron transfer. The resulting dinuclear Pd species **5**, featuring a monothiolate bridgehead, contains a unique mixed-valence ligand set, with one PNO^{ISO} and one PNO^{AP} unit. The introduction of a flanking phosphine group could allow the expansion of the concept of ligand-induced electron transfer and radical-type reactivity to “softer” low-valent noble metals.

Received: October 13, 2014

Revised: October 30, 2014

Published online: November 27, 2014

Keywords: bond activation · mixed valency · palladium · phosphane ligands · redox-active ligands

- [1] a) W. Kaim, B. Schwederski, *Coord. Chem. Rev.* **2010**, *254*, 1580–1588; b) L. Que, W. B. Tolman, *Nature* **2008**, *455*, 333–340; c) D. L. Harris, *Curr. Opin. Chem. Biol.* **2001**, *5*, 724–735.
- [2] a) C. G. Pierpont, *Coord. Chem. Rev.* **2001**, *219*, 415–433; b) P. Zanello, M. Corsini, *Coord. Chem. Rev.* **2006**, *250*, 2000–2022; c) K. Ray, T. Petrenko, K. Wieghardt, F. Neese, *Dalton Trans.* **2007**, 1552–1566; d) R. Eisenberg, H. B. Gray, *Inorg. Chem.* **2011**, *50*, 9741–9751.
- [3] a) J. I. van der Vlugt, *Eur. J. Inorg. Chem.* **2012**, 363–375; b) V. Lyaskovskyy, B. de Bruin, *ACS Catal.* **2012**, *2*, 270–279; c) A. L. Smith, K. I. Hardcastle, J. D. Soper, *J. Am. Chem. Soc.* **2010**, *132*, 14358–14360; d) K. J. Sylvester, P. J. Chirik, *J. Am. Chem. Soc.* **2009**, *131*, 8772–8774; e) M. R. Ringenberg, S. L. Kokatam, Z. M. Heiden, T. B. Rauchfuss, *J. Am. Chem. Soc.* **2008**, *130*, 788–789; f) M. Königsmann, N. Donati, D. Stein, H. Schönberg, J. Harmer, A. Sreekanth, H. Grützmacher, *Angew. Chem. Int. Ed.* **2007**, *46*, 3567–3570; *Angew. Chem.* **2007**, *119*, 3637–3640.
- [4] a) K. G. Caulton, *Eur. J. Inorg. Chem.* **2012**, 435–443; b) W. Kaim, *Inorg. Chem.* **2011**, *50*, 9752–9765.
- [5] A. I. Poddelsky, V. K. Cherkasov, G. A. Abakumov, *Coord. Chem. Rev.* **2009**, *253*, 291–324.
- [6] a) N. P. Mankad, E. Rivard, S. B. Harkins, J. C. Peters, *J. Am. Chem. Soc.* **2005**, *127*, 16032–16033; b) Y. M. Rhee, M. Head-Gordon, *J. Am. Chem. Soc.* **2008**, *130*, 3878–3887; c) V. Lyaskovskyy, R. J. A. van Dijk-Moes, S. Burck, W. I. Dzik, M. Lutz, A. W. Ehlers, J. C. Slootweg, B. de Bruin, K. Lammertsma, *Organometallics* **2013**, *32*, 363–373.
- [7] Organic P-based radicals: a) S. Loss, A. Magistrato, L. Cataldo, S. Hoffmann, M. Geoffroy, U. Röthlisberger, H. Grützmacher, *Angew. Chem. Int. Ed.* **2001**, *40*, 723–726; *Angew. Chem.* **2001**, *113*, 749–751; b) S. Marque, P. Tordo, *Top. Curr. Chem.* **2005**, *250*, 43–76; c) O. Back, B. Donnadieu, P. Parameswaran, G. Frenking, G. Bertrand, *Nat. Chem.* **2010**, *2*, 369–373; d) S. Ishida, F. Hirakawa, T. Iwamoto, *J. Am. Chem. Soc.* **2011**, *133*, 12968–12971; e) N. A. Giffin, A. D. Hendsbee, T. L. Roemmele, M. D. Lumsden, C. C. Pye, J. D. Masuda, *Inorg. Chem.* **2012**, *51*, 11837–11850.
- [8] M. W. Bezpalko, B. M. Foxman, C. M. Thomas, *Inorg. Chem.* **2013**, *52*, 12329–12331.
- [9] Metallocenes and tetrathiofulvenes: a) G. Bandoli, A. Dolmella, *Coord. Chem. Rev.* **2000**, *209*, 161–196; b) N. Avarvari, M. Fourmigué, *Chem. Commun.* **2004**, 1300–1301; Diphenylamido-phosphines: c) D. Adhikari, S. Mossin, F. Basuli, J. C. Huffman, R. K. Szilagy, K. Meyer, D. J. Mindiola, *J. Am. Chem. Soc.* **2008**, *130*, 3676–3682; d) S. B. Harkins, N. P. Mankad, A. J. M. Miller, R. K. Szilagy, J. C. Peters, *J. Am. Chem. Soc.* **2008**, *130*, 3478–3485; e) R. Lindner, B. van den Bosch, M. Lutz, J. N. H. Reek, J. I. van der Vlugt, *Organometallics* **2011**, *30*, 499–510.
- [10] Recently, Hicks and co-workers reported coordination of a redox-active verdazyl phosphine to Pd^{II}: C. A. Sanz, M. J. Ferguson, R. McDonald, B. O. Patrick, R. G. Hicks, *Chem. Commun.* **2014**, *50*, 11676–11678.
- [11] Reductive C–C coupling: J. M. Darmon, S. C. E. Stieber, K. T. Sylvester, I. Fernández, E. Lobkovsky, S. P. Semproni, E. Bill, K. Wieghardt, S. DeBeer, P. J. Chirik, *J. Am. Chem. Soc.* **2012**, *134*, 17125–17137.
- [12] Cleavage of disulfides: I. L. Fedushkin, A. S. Nikipelov, A. A. Skatova, O. V. Maslova, A. N. Lukoyanov, G. K. Fukin, A. V. Cherkasov, *Eur. J. Inorg. Chem.* **2009**, 3742–3749.
- [13] Reductive thiolate coupling: J. L. Wong, R. H. Sánchez, J. G. Logan, R. A. Zarkesh, J. W. Ziller, A. F. Heyduk, *Chem. Sci.* **2013**, *4*, 1906–1910.
- [14] Dioxygen activation: C. A. Lippert, S. A. Arnstein, C. D. Sherill, J. D. Soper, *J. Am. Chem. Soc.* **2010**, *132*, 3879–3892; Note: this overall two-electron process is proposed to proceed through two sequential ligand-to-substrate one-electron transfer steps.
- [15] a) C. Díaz, C. Leal, N. Yutronic, *J. Organomet. Chem.* **1996**, *516*, 59–64; b) B. L. Tran, C.-H. Chen, D. J. Mindiola, *Inorg. Chim. Acta* **2011**, *369*, 215–222; c) A. O. Tolpygin, A. V. Cherkasov, G. K. Fukin, A. A. Trifonov, *Inorg. Chem.* **2014**, *53*, 1537–1543; d) P. Pelties, D. Herrmann, B. de Bruin, F. Hartl, R. Wolf, *Chem. Commun.* **2014**, *50*, 7014–7016.
- [16] Radicals in biology: a) J. Stubbe, W. A. van der Donk, *Chem. Rev.* **1998**, *98*, 705–762; b) J. W. Whittaker, *Chem. Rev.* **2003**, *103*, 2347–2363; Radicals in synthesis: c) W. Buckel, *Angew. Chem. Int. Ed.* **2009**, *48*, 6779–6787; *Angew. Chem.* **2009**, *121*, 6911–6920; d) J. Hioe, H. Zipse, *Org. Biomol. Chem.* **2010**, *8*, 3609–3617; e) W. I. Dzik, X. P. Zhang, B. de Bruin, *Inorg. Chem.* **2011**, *50*, 9896–9903; f) M. Heinrich, A. Gansäuer, *Top. Curr. Chem.* **2012**, *320* (Radicals in Organic Synthesis III), Springer; g) F. Déniš, M. Pichowicz, G. Povie, P. Renaud, *Chem. Rev.* **2014**, *114*, 2587–2693.
- [17] D. L. J. Broere, B. de Bruin, J. N. H. Reek, M. Lutz, S. Dechert, J. I. van der Vlugt, *J. Am. Chem. Soc.* **2014**, *136*, 11574–11577.
- [18] Condensation between *o*-iodoaniline and 3,5-di-*tert*-butylcatechol followed by Pd-catalyzed coupling with HPPH₂. Modified procedure from: S. Wanniarachchi, J. S. Hewage, S. V. Lindeman, J. R. Gardinier, *Organometallics* **2013**, *32*, 2885–2888.
- [19] a) O. M. Ní Dhubhghaill, J. Lennon, M. G. B. Drew, *Dalton Trans.* **2005**, 3213–3220; b) B. Crociani, S. Antonaroli, M. Burattini, P. Paoli, P. Rossi, *Dalton Trans.* **2010**, *39*, 3665–3672.
- [20] a) J. C. Peters, S. B. Harkins, S. D. Brown, M. W. Day, *Inorg. Chem.* **2001**, *40*, 5083–5091; b) X. Sun, H. Chun, K. Hildenbrand, E. Bothe, T. Weyhermüller, F. Neese, K. Wieghardt, *Inorg. Chem.* **2002**, *41*, 4295–4303; c) K. S. Min, T. Weyhermüller, E. Bothe, K. Wieghardt, *Inorg. Chem.* **2004**, *43*, 2922–2931; d) K. Chlopek, E. Bothe, F. Neese, T. Weyhermüller, K. Wieghardt, *Inorg. Chem.* **2006**, *45*, 6298–6307; e) N. Gómez-Blanco, J. J. Fernández, A. Fernández, D. Vázquez-García, M. López-Torres, J. M. Vila, *Eur. J. Inorg. Chem.* **2009**, 3071–3083.

- [21] The simulated spectrum and calculated hyperfine couplings correlate well with the experimental data, with the exception of some hyperfine interactions (see the Supporting Information).
- [22] a) D. Shimizu, N. Takeda, N. Tokitoh, *J. Organomet. Chem.* **2007**, *692*, 2716–2728; b) P. K. Dutta, A. K. Asatkar, S. S. Zade, S. Panda, *Dalton Trans.* **2014**, *43*, 1736–1743.
- [23] Proton-transfer reaction of thiols with Pd^I: a) S. J. Ling Foo, N. D. Jones, B. O. Patrick, B. R. James, *Chem. Commun.* **2003**, 988–989; b) D. P. Hruszkewycz, J. Wu, N. Hazari, C. D. Incarvito, *J. Am. Chem. Soc.* **2011**, *133*, 3280–3283; With Pd^{II}: c) J. Ruiz, N. Cutillas, J. Torregrosa, G. Garcia, G. Lopez, P. A. Chaloner, P. B. Hitchcock, R. M. Harrison, *J. Chem. Soc. Dalton Trans.* **1994**, 2353–2357; Metathesis with TISR: d) R. Usón, M. A. Usón, S. Herrero, L. Rello, *Inorg. Chem.* **1998**, *37*, 4473–4476.
- [24] Ligand mixed valency induced through selective one-electron ligand-based redox in dinuclear complexes with two bridging ligands has been reported: a) S. Mukherjee, T. Weyhermüller, E. Bothe, K. Wieghardt, P. Chaudhuri, *Dalton Trans.* **2004**, 3842–3853; b) N. Roy, S. Sproules, T. Weyhermüller, K. Wieghardt, *Inorg. Chem.* **2009**, *48*, 3783–3791; For an example of solid state localized ligand mixed valency, see: c) S. Kokatam, T. Weyhermüller, E. Bothe, P. Chaudhuri, K. Wieghardt, *Inorg. Chem.* **2005**, *44*, 3709–3717.
- [25] a) J. A. Thomas, *Coord. Chem. Rev.* **2013**, *257*, 1555–1563; b) M. G. Sommer, D. Schweinfurth, F. Weisser, S. Hohloch, B. Sarkar, *Organometallics* **2013**, *32*, 2069–2078; c) S. D. J. McKinnon, B. O. Patrick, A. B. P. Lever, R. G. Hicks, *Inorg. Chem.* **2013**, *52*, 8053–8066; d) M. A. Fox, B. Le Guennic, R. L. Roberts, D. A. Brue, D. S. Yufit, J. A. K. Howard, G. Manca, J.-F. Halet, F. Hartl, P. J. Low, *J. Am. Chem. Soc.* **2011**, *133*, 18433–18446; e) W. Kaim, G. K. Lahiri, *Angew. Chem. Int. Ed.* **2007**, *46*, 1778–1796; *Angew. Chem.* **2007**, *119*, 1808–1828.
- [26] a) V. Lloveras, J. Vidal-Gancedo, D. Ruiz-Molina, T. M. Figueira-Duarte, J. F. Nierengarten, J. Veciana, C. Rovira, *Faraday Discuss.* **2006**, *131*, 291–305; b) V. Lloveras, J. Vidal-Gancedo, T. M. Figueira-Duarte, J. F. Nierengarten, J. J. Novoa, F. Mota, N. Ventosa, J. Veciana, C. Rovira, *J. Am. Chem. Soc.* **2011**, *133*, 5818–5833.
- [27] Reduction potential vs. SCE: PhSSPh –1.6 V; *t*BuSS*t*Bu –2.71 V. Bond dissociation energy (BDE) of PhSSPh 55.0 kcal mol⁻¹; dialkyldisulfides ≈ 65 kcal mol⁻¹; MeSSMe: 73.2 kcal mol⁻¹.
- [28] a) see ref. [16g]; b) O. Ito, M. Matsuda, *J. Am. Chem. Soc.* **1983**, *105*, 1937–1940.